

Reaching the continuum limit in lattice gauge theory - without a computer.

John A.L. McIntosh and Lloyd C.L. Hollenberg

Research Centre for High Energy Physics

School of Physics

The University of Melbourne

Victoria 3010, Australia

(j.mcintosh@physics.unimelb.edu.au, l.hollenberg@physics.unimelb.edu.au)

The scaling slope of the anti-symmetric mass gap M of compact $U(1)_{2+1}$ lattice gauge theory is obtained analytically in the Hamiltonian formalism using the plaquette expansion. Based on the first four moments of the Hamiltonian with respect to a one-plaquette mean field state the results demonstrate clear scaling of M at and beyond the transition from strong to weak coupling. The scaling parameters determined agree well with the range of numerical determinations available.

PACS number: 11.15.Ha

With the vast majority of calculations carried out in lattice gauge theory being of an intensely numerical nature, non-perturbative analytic results are of considerable interest. If we look to complex theories such as QCD, the most basic quantities one wishes to calculate are the states of the mass spectrum of the theory. In lattice gauge theory, such non-perturbative calculations are carried out numerically in the path integral representation to some level of approximation by evaluating the correlation functions by Monte-Carlo simulation. Typically, one must work with large lattices and spacings which are fine enough to detect the continuum behaviour of the theory in question. The computational resources employed in this approach are enormous, essentially reflecting the seemingly intractable many-body problem at the core of non-perturbative quantum field theory. We report here on a novel approach to studying lattice gauge theories, and as an indication of its potential apply the method to compact $U(1)$ lattice gauge theory in $2+1$ dimensions. This theory, although obviously simpler than QCD itself, nevertheless shares many of the features of the QCD many-body problem on the lattice, namely similar numerical difficulties in calculating masses in the continuum limit. The scaling behaviour for the mass gap, M , is expected [1] to be:

$$M^2 a^2 \equiv \beta e^{-k_0 \beta + k_1}, \quad (1)$$

where k_0 and k_1 are constants, a is the lattice spacing and $\beta = 1/g^2$ where g is the dimensionless coupling constant. As with most other interesting lattice gauge theories, it has long been a maxim that even $U(1)_{2+1}$ lattice gauge theory must be studied by large scale numerical computation in order to determine the lowest mass state of the theory. Indeed, although studied by a range of different methods, the scaling properties of this theory are still not known precisely: $k_0 \sim 4.1 - 6.345$ and $k_1 \sim 4.369 - 6.27$ (see [2] for summary). In this sense the results we present are quite spectacular – an analytic determination of the mass-gap, which clearly demonstrates the scaling behaviour expected in the continuum limit with $k_0 = 4.42$ and $k_1 = 5.41$.

Throughout this paper we will work in the Hamiltonian formalism. When one speaks of “lattice gauge theory” it is usually meant the Lagrangian formalism where one essentially converts the theory to a statistical model in Euclidean space into order to facilitate the use of powerful Monte-Carlo methods. In contrast, there is a dearth of equivalently efficient methodology in the Hamiltonian formalism of lattice gauge theory. This fact has been responsible for the domination of action-based lattice gauge theory over the Hamiltonian approach. While there exist numerical and Monte-Carlo methods applicable to the Hamiltonian lattice formalism, these have been of limited use, and certainly the results are not of the same quality as in the action-based approach. However, there is a certain appeal of the Hamiltonian formalism which tugs at physicists’ intuition. There are also technical reasons for considering the Hamiltonian approach – one often works directly on infinite lattices, fermions are simpler to incorporate (no fermion determinant), the lattices are 3D as opposed to 4D (time is continuous), and finite temperature physics may actually be more natural in this formalism [3]. The lack of powerful and systematic methodology, such as is available in the Lagrangian formalism, has prevented these features from being exploited or investigated – hence, the importance of new methodologies. Here we use a novel approach – the plaquette expansion – based on a large volume expansion of Lanczos tridiagonalization.

The Kogut-Susskind Hamiltonian for $U(1)_{2+1}$ on a lattice of N plaquettes is given by [4]:

$$H = \frac{g^2}{2} \sum_l \hat{E}_l^2 + \frac{1}{2g^2} \sum_p \left[2 - (\hat{U}_p + \hat{U}_p^\dagger) \right], \quad (2)$$

The strong-coupling limit is defined by $g \rightarrow \infty$ and the weak-coupling limit by $g \rightarrow 0$. For the $U(1)$ model, $\hat{E}_l$ is the electric field operator associated with lattice link l . In the representation in which $\hat{E}_l$ is diagonal, the eigenvalue E_l takes on integer values $0, \pm 1, \pm 2, \dots$ on each link and the link operators $\hat{U}_l$ and $\hat{U}_l^\dagger$ raise or lower the

value of E_l on link l by unity respectively due to the commutation relations:

$$[\hat{E}_l, \hat{U}_l] = \hat{U}_l \quad , \quad [\hat{E}_l, \hat{U}_l^\dagger] = -\hat{U}_l^\dagger . \quad (3)$$

The operator $\hat{U}_p$ acts on the links around the smallest closed (Wilson) loop or square on the lattice known commonly as a plaquette p :

$$\hat{U}_p = \hat{U}_1 \hat{U}_2 \hat{U}_3^\dagger \hat{U}_4^\dagger . \quad (4)$$

The extensive nature of the Hamiltonian Equation (2) lends itself to the application of the plaquette expansion. Indeed this method was originally devised to study such local lattice models [5]. Beginning with the well known Lanczos recurrence for the iterative transformation of the Hamiltonian into tri-diagonal form, a trial state $|\psi_1\rangle$ with the desired symmetries of the state of interest $|\Psi_0\rangle$ (the ground state or vacuum in this case) and non-zero overlap (*i.e.* $\langle\psi_1|\Psi_0\rangle \neq 0$) is chosen and a basis $\{|\psi_n\rangle\}$ is constructed according to

$$|\psi_n\rangle = \frac{1}{\beta_{n-1}} [(H - \alpha_{n-1})|\psi_{n-1}\rangle - \beta_{n-2}|\psi_{n-2}\rangle] , \quad (5)$$

where $\alpha_n = \langle\psi_n|H|\psi_n\rangle$ and $\beta_n = \langle\psi_{n+1}|H|\psi_n\rangle$ are the matrix elements of the Hamiltonian in tri-diagonal form.

The matrix elements, α_n and β_n , can be written in terms of Hamiltonian moments $\langle H^n \rangle \equiv \langle\psi_1|H^n|\psi_1\rangle$. As the connected part of the Hamiltonian moment is proportional to the volume of the system $\langle H^n \rangle_c = e_n N$, one may utilize the cumulant transformation [6],

$$\langle H^n \rangle_c = \langle H^n \rangle - \sum_{p=0}^{n-2} \binom{n-1}{p} \langle H^{p+1} \rangle_c \langle H^{n-1-p} \rangle , \quad (6)$$

and rewrite the α_n and β_n in terms of the connected vacuum coefficients e_n . Extensivity of the problem leads to the following plaquette expansions in $1/N$ [5]:

$$\frac{\alpha_n}{N} = e_1 + s \left[\frac{e_3}{e_2} \right] + O(s^2) \equiv \bar{\alpha}(s), \quad (7)$$

$$\frac{\beta_n^2}{N^2} = s e_2 + \frac{1}{2} s^2 \left[\frac{e_2 e_4 - e_3^2}{e_2^2} \right] + O(s^3) \equiv \bar{\beta}^2(s), \quad (8)$$

where $s \equiv n/N$. In the bulk limit ($n, N \rightarrow \infty$), keeping s fixed, it was shown in [7] that one may perform the exact diagonalization of the Lanczos tri-diagonal matrix for the vacuum energy density ε_0 :

$$\varepsilon_0 = \inf (\bar{\alpha}(s) - 2\bar{\beta}(s)) \quad (9)$$

For low order truncations of the series (employing moments up to $\langle H^r \rangle$) one can minimize (9) with respect to s analytically and obtain the closed form expressions for the ground state energy, $\varepsilon_0(r)$:

$$\varepsilon_0(1) = e_1,$$

$$\begin{aligned} \varepsilon_0(3) &= e_1 - \frac{e_2^2}{e_3}, \\ \varepsilon_0(4) &= e_1 + \frac{e_2^2}{e_2 e_4 - e_3^2} \left[\sqrt{3e_3^2 - 2e_2 e_4 - e_3} \right]. \end{aligned} \quad (10)$$

Note the first approximant $\varepsilon_0(1)$ is just the variational result. The expressions for $\varepsilon_0(3)$ and $\varepsilon_0(4)$ clearly illustrate the non-perturbative nature of the plaquette expansion as an effective summation over clusters of increasing complexity.

To determine expressions for excited states we note that lowest state of a given sector, S , in the spectrum will in general have an energy:

$$E_0^{(S)} = \varepsilon_0 N + M_0^{(S)} . \quad (11)$$

where $M_0^{(S)}$ is the mass gap. For excited states the Hamiltonian moments have the form $\langle H^n \rangle_c^{(S)} = e_n N + m_n^{(S)}$. Thus, we can then define analogous ‘‘cumulants’’ $e_n^{(S)}$ for this sector of Hilbert space as $e_n^{(S)} \equiv e_n + m_n^{(S)}/N$, where we call $m_n^{(S)}$ the mass gap connected moments. One can derive expressions for approximants, $M_0^{(S)}(r)$, to the mass gap, in a similar fashion to those for the vacuum energy density. For example, using up to 4th order moments we have [8]:

$$M_0^{(S)}(4) = \sum_{n=1}^4 m_n^{(s)} e_2^{n-1} F_n, \quad (12)$$

where the vacuum moment functions, F_n , are given by

$$\begin{aligned} F_1 &= 1, \\ F_2 &= \frac{\sqrt{\Delta} (\Omega) + e_3 (-\Omega + e_2 e_3^2 e_4 - e_2^2 e_4^2)}{(\Delta) (e_3^2 - e_2 e_4)^2}, \\ F_3 &= \frac{(2e_2^2 e_4^2 - 3e_3^4 - e_2 e_3^2 e_4) + e_3 \sqrt{\Delta} (3e_3^2 - e_2 e_4)}{(\Delta) (e_3^2 - e_2 e_4)^2}, \\ F_4 &= \frac{\sqrt{\Delta} (e_2 e_4 - 2e_3^2) + e_3 (\Delta)}{(\Delta) (e_3^2 - e_2 e_4)^2}, \end{aligned} \quad (13)$$

with $\Delta = 3e_3^2 - 2e_2 e_4$ and $\Omega = 8e_2 e_3^2 e_4 - 6e_3^4 - 3e_2^2 e_4^2$.

It is appropriate to comment on relevant methods, which by and large employ trial states encompassing as much of the expected continuum physics as is viable. Often, such calculations have employed the simplest gauge invariant trial state — the strong-coupling vacuum $|0\rangle$ (defined as the state satisfying $\hat{E}_l|0\rangle = 0$ for every link). This is the perturbative starting point for series calculations, which are then extrapolated to weak coupling. Although the calculation of moments with respect to this state can be carried out to relatively high order, typically, one finds that this state is simply inadequate to explore the non-perturbative weak-coupling regime of the theory. In the case of the plaquette expansion, a window of

scaling in terms of the expansion order was evident, but higher order results became problematic [9].

More recently, trial states which are more complex and appropriate to weak-coupling have been investigated by a number of authors [10–15] in varying contexts. There is a clear tradeoff between physical complexity of the trial state, and algebraic complexity of the moment calculation. Given this, we utilized a one plaquette exponential trial state, that has the correct weak coupling behaviour for the vacuum energy density – the so-called *mean-field* state – defined as: $|\psi_1\rangle = \exp\{\hat{S}/2\}|0\rangle$, where $\hat{S} = \lambda \sum_p \{\hat{U}_p + \hat{U}_p^\dagger\}$ and λ will eventually be determined variationally from $\langle H \rangle$. It is convenient to work in the so-called U representation where the trial state is a function of U_p variables:

$$\psi_1[U_p] = e^{S[U_p]/2} = \exp \left\{ \lambda \sum_p \{U_p + U_p^\dagger\} \right\} \quad (14)$$

As the state is constructed from plaquette variables, U_p , it is automatically gauge invariant. One expects the true ground state of the system to have contributions from all possible Wilson loops – an important feature of the plaquette expansion method is that by its very nature larger sized Wilson loops (larger clusters) are systematically introduced.

In $D = 2 + 1$ the Hamiltonian moments can be written in terms of integrals over plaquette variables, for example the vacuum moments are given schematically by:

$$\langle H^n \rangle = \frac{\int \mathcal{D}U_p \psi_1[U_p]^\dagger H^n \psi_1[U_p]}{\int \mathcal{D}U_p \psi_1[U_p]^\dagger \psi_1[U_p]} \equiv \frac{\int \mathcal{D}U_p G_n[U_p] e^{S[U_p]}}{\int \mathcal{D}U_p e^{S[U_p]}}$$

For the group $U(1)$, the plaquette variables may be rewritten in terms of plaquette phase angles $U_p \equiv \exp(i\phi_p)$. The calculation of Hamiltonian moments for use in the plaquette expansion utilizing the one-plaquette state in $2 + 1$ dimensions comes down to evaluating integrals of the following form,

$$\langle H^n \rangle \equiv \left[\frac{1}{2\pi I_0(2\lambda)} \right]^N \int_{-\pi}^{\pi} \prod_p d\phi_p G_n[\phi_p] e^{S[\phi_p]}, \quad (15)$$

where $I_0(2\lambda)$ is the zeroth-order modified Bessel function of the first kind. The integrand functions, $G_n[\phi_p]$, naturally become more complex as one increases the moment order, however, all integrations can be carried out analytically. All expressions are analytic in λ and β , and once $\langle H \rangle$ is minimized we implicitly determine $\lambda(\beta)$.

It is well known [12,16] that the variational calculation $\langle H \rangle$ applying the one-plaquette state in $D = 2 + 1$ dimensions involves no correlations between plaquettes and thus one is simply solving a Mathieu-type problem. In this context, the benefit of choosing the one-plaquette state as the trial state rather than the strong-coupling vacuum becomes more apparent on investigation of the

strong ($\beta \rightarrow 0$) and weak-coupling ($\beta \rightarrow \infty$) limits of $\langle H \rangle$. For the strong-coupling limit, one finds for the energy density: $\varepsilon_0(1) = \langle H \rangle / N \sim \beta - \beta^3/4 + O(\beta^7)$, which agrees with strong-coupling perturbation theory [17]. Similarly, for the weak-coupling limit, one finds: $\varepsilon_0(1) \sim A - A^2/(8\beta) + O(1/\beta^2)$, with $A = 1$. This agrees exactly with weak-coupling perturbation theory for the Mathieu problem to the order shown. Exact $U(1)_{2+1}$ has a similar weak-coupling expansion [18,19] with $A = 0.9581$. With the inclusion of larger sized Wilson loops obviously the $U(1)$ calculations, as they should, will deviate from the Mathieu results. As this trial state encompasses the actual behaviour in both limits, one has confidence in using the plaquette expansion to systematically obtain results throughout the range of β .




FIG. 1. Mass gap results applying one (minimized numerically), three and four one-plaquette Hamiltonian moments. Scaling is evident for the four order moments curve from $\beta = 0.7$ to 1.5 .

Using a diagrammatic method (details to be presented elsewhere) we were able to derive the first four one-plaquette connected moments for both the vacuum and anti-symmetric state. These moments immediately give analytic approximates for vacuum energy and the mass gap.

Since the expected scaling behaviour for compact $U(1)_{2+1}$ is of the form (1), the one-plaquette state results for the plaquette expansion to first order are presented as a $\ln(M^2/\beta)$ versus β plot in Figure 1. The results using one, three and four moments appear to be converging towards the scaling slope. Clearly scaling is evident for inverse-coupling values $\beta = 0.7$ to 1.5 , passing the transition point at $\beta = 1$. The scaling form we find is given by $M^2 = \beta \exp(-4.34\beta + 5.34)$, and agrees well with other estimates (as summarized in [2]).

It should be noted that the minimization of $\langle H \rangle$ is the only strictly numerical procedure (small as it is) in our calculation. A totally analytic estimation (keeping in line with the title of this paper) is possible when one writes the function $\lambda(\beta)$ which minimizes $\langle H \rangle$ as a series expansion

sion: $\lambda = \beta^2/2 - \beta^6/16 + \dots$. Keeping the first five terms of the expansion (for which the resulting values have converged), scaling is observed from $\beta = 0.8$ to 1.2 (Figure 2). The series breaks down at larger β values. The scaling form is given by $M^2 = \beta \exp(-4.42\beta + 5.41)$, and again agrees well with the numerical estimates available.

In summary, applying the plaquette expansion we were able to identify scaling of the anti-symmetric (or photon) mass gap in the weak-coupling regime of compact $U(1)_{2+1}$ lattice gauge theory via a completely analytic calculation utilizing only four one-plaquette state Hamiltonian moments. Furthermore, although we have presented results for only the first four moments (for which the calculation is entirely analytic) it should in principle be possible to derive higher moments with respect to this state. Indeed we have preliminary results for the vacuum energy density to sixth order [2] giving high accuracy over a large range of couplings.




FIG. 2. Convergence of the complete analytic mass gap results for one (variational), three and four one-plaquette Hamiltonian moments to the continuum limit, using the series form for $\lambda(\beta)$.

These results may have important implications for lattice gauge theory – the plaquette expansion method itself can be applied to any extensive many-body problem. The only existing application of the method to QCD_{3+1} (including massless quarks) with respect to the strong coupling state proved somewhat disappointing [20] in that scaling of the nucleon/meson mass ratio was not evident. However, the results presented here indicate that the one-plaquette state may be of sufficient quality to capture the essence of the continuum physics in lattice gauge theory at relatively low orders. An exciting possibility is that the continuum limit might finally be observed in the Hamiltonian formalism for QCD using this method.

To finish, we propose a semi-analytic program to study the 3+1 lattice gauge theories in the Hamiltonian formalism. The main obstacle to overcome is the calculation of the moments in 3+1 dimensions. The appeal of 2+1

systems as test models is that the transformation from link variables to plaquette variables is trivial and makes the integrations tractable – hence the analytic work reported in this paper. In 3+1 dimensions it is well known that the transformation cannot be carried out in closed form due to the appearance of Biachi identities. However, the calculation of moments in 3+1 dimensions by Monte-Carlo methods is actually a relatively small scale numerical exercise. Because we are dealing with a cluster expansion (we need only connected moments) the lattices required fairly modest in extent, and are by definition only three dimensional. The integrands become complicated as larger correlations are included, which means that the statistics must be very good, however, preliminary calculations have demonstrated that the required precision is possible to achieve with quite modest computing resources [2].

This work was supported by the Australian Research Council and the Australian Postgraduate Award scheme.

-
- [1] M.Göpfert and G.Mack, *Commun. Math. Phys.* **82**, 545 (1982).
 - [2] J.A.L.McIntosh, Analytic Lanczos Methodology and Scaling in Hamiltonian Lattice Gauge Theory, Ph.D. thesis (2000).
 - [3] E.B.Gregory, S.H.Guo, H.Kröger and X.Q.Luo, *Phys. Rev. D* **62**, 054508 (2000).
 - [4] J.Kogut and L.Susskind, *Phys. Rev. D* **11**, 395 (1975).
 - [5] L.C.L.Hollenberg, *Phys. Rev. D* **47**, 1640 (1993).
 - [6] D.Horn and M.Weinstein, *Phys. Rev. D* **30**, 1256 (1984).
 - [7] L.C.L.Hollenberg and N.S.Witte, *Phys. Rev. B* **54**, 16309 (1996).
 - [8] L.C.L.Hollenberg, M.Wilson and N.S.Witte, *Phys. Lett. B* **361**, 81 (1995).
 - [9] J.A.L.McIntosh and L.C.L.Hollenberg, *Z. Phys. C* **76**, 175 (1997).
 - [10] W.Langguth, *Z.Phys. C* **23**, 289 (1984).
 - [11] D.W.Heys and D.R.Stump, *Nucl. Phys. B* **257**, 19 (1985).
 - [12] S.J.Baker, R.F.Bishop and N.J.Davidson, *Phys. Rev. D* **53**, 2610 (1996).
 - [13] S.J.Baker, R.F.Bishop and N.J.Davidson, *Nucl. Phys. B (Proc. Suppl.)* **53**, 834 (1997).
 - [14] A.Duncan and R.Roskies, *Phys. Rev. D* **31**, 364 (1985).
 - [15] A.Duncan, *Nucl. Phys. B* **258**, 125 (1985).
 - [16] R.F.Bishop, A.S.Kendall, L.Y.Wong and Y.Xian, *Phys. Rev. D* **48**, 887 (1993).
 - [17] A.C.Irving and C.J.Hamer, *Nucl. Phys. B* **230**, 361 (1984); *Nucl. Phys. B* **235**, 358 (1984).
 - [18] A.Dabringhaus, M.L.Ristig and J.W.Clark, *Phys. Rev. D* **43**, 1978 (1991).
 - [19] C.J.Hamer and Z.Weihong, *Phys. Rev. D* **48**, 4435 (1993).
 - [20] L.C.L.Hollenberg, *Phys. Rev. D* **50**, 6917 (1994).